



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?

Citation for published version:

Popkirov, S, Carson, AJ & Stone, J 2018, 'Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?', *Seizure - European Journal of Epilepsy*.
<https://doi.org/10.1016/j.seizure.2018.04.009>

Digital Object Identifier (DOI):

[10.1016/j.seizure.2018.04.009](https://doi.org/10.1016/j.seizure.2018.04.009)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Seizure - European Journal of Epilepsy

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Accepted Manuscript

Title: Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?

Authors: Stoyan Popkirov, Alan J. Carson, Jon Stone

PII: S1059-1311(18)30109-2
DOI: <https://doi.org/10.1016/j.seizure.2018.04.009>
Reference: YSEIZ 3164

To appear in: *Seizure*

Received date: 13-2-2018
Revised date: 31-3-2018
Accepted date: 10-4-2018



Please cite this article as: Popkirov Stoyan, Carson Alan J, Stone Jon. Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?. *SEIZURE: European Journal of Epilepsy* <https://doi.org/10.1016/j.seizure.2018.04.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

REVISION 1

Ref: SEIZURE_2018_90

Title:

Scared or scarred: could ‘dissociogenic’ lesions predispose to nonepileptic seizures after head trauma?

Category:

Focused review

Authors:

Stoyan Popkirov^{a*}, Alan J. Carson^{b,c}, Jon Stone^b

Affiliations:

^a Department of Neurology, University Hospital Knappschaftskrankenhaus Bochum, Ruhr-University Bochum, Bochum, Germany

^b Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, United Kingdom

^c Scottish Neurobehavioural Rehabilitation Unit, Royal Edinburgh Hospital, Edinburgh, United Kingdom

*Corresponding author:

Dr. Stoyan Popkirov, Klinik für Neurologie, Universitätsklinikum Knappschaftskrankenhaus Bochum, In der Schornau 23-25, 44892 Bochum, Germany.

Email: popkirov@gmail.com.

Phone: +49 234 299-3701.

Fax: +49 234 299-3719

Highlights

- Traumatic brain injury (TBI) is common in psychogenic nonepileptic seizures (PNES).
- A possible connection is seen in maladaptive symptom modelling.
- Effects of mild brain injury can include disruption of functional connectivity.
- This disruption might temporarily facilitate dissociation and PNES development.
- Mild TBI might be a cause of “dissociogenic” lesions in some patients.

Abstract

A history of head injury is common in patients with psychogenic nonepileptic seizures (PNES). This association has so far been interpreted as either spurious or psychologically mediated. Biased recall and misattribution could foster illness beliefs about brain damage that promote symptom production. Furthermore, the emotional impact of head injury could induce long-term changes in stress responsivity. Lastly, maladaptive cognitive-behavioural processes involving symptom modelling and aversive conditioning, known to play a role in functional neurological disorders, could contribute to the development of PNES after head trauma. Lesional effects of head injury, on the other hand, remain unexplored in the context of PNES. However, even mild traumatic brain injury without structural MRI abnormalities on routine imaging can lead to disruptions of network connectivity that correlate with short-term cognitive impairments and psychiatric symptoms. Since alterations in global functional connectivity have been demonstrated in PNES patients using imaging and electroencephalography, we hypothesize that, in some patients, TBI and the associated disruption of long-range association fibres could contribute to the individual propensity for dissociative experiences in general and PNES in particular. This possibility is explored in the context of new cognitive-behavioural models of PNES pathogenesis, and the concept of a "dissociogenic" brain lesion is introduced.

Keywords

traumatic brain injury; concussion; psychogenic nonepileptic seizures; dissociative seizures; dissociation; functional connectivity

Introduction

Psychogenic nonepileptic seizures (PNES) are paroxysmal episodes of dissociation that can superficially resemble epileptic seizures or syncope, but do not arise from neuronal hypersynchronisation or ischaemia.¹ They can occur in isolation as a form of functional neurological (dissociative) disorder, or in association with various neurological and psychiatric conditions such as epilepsy or posttraumatic stress disorder (PTSD). The underlying psychophysiological processes are still a matter of controversial debate, and closely associated constructs such as somatization, conversion and dissociation remain themselves inconsistently defined (see [1] for review). Recently accelerated efforts to elucidate the neural ("biological") underpinnings of PNES have shed some light on specific physiological dysfunctions and anatomical predispositions for PNES, challenging a dualistic view of a disorder often seen as purely psychogenic.^{1,2} Using the common association of traumatic

brain injury (TBI) and PNES as a starting point, this article aims to expand our understanding of PNES pathophysiology with the proposition of “dissociogenic” brain lesions. In analogy to epileptogenic lesions, which can (but do not always) cause focal epileptic seizures, it will be argued that dissociogenic lesions can predispose individuals to PNES in addition to known risk factors such as childhood trauma, anxiety or personality disorder.

TBI preceding PNES

Early case note reviews from epilepsy centres demonstrated an unexpectedly high comorbidity of PNES and TBI.^{3,4} Studies have since then confirmed this observation with rates of co-occurrence varying between 16-83%, and a weighted average of 43% among 1039 adult PNES patients across 17 studies (Table 1).³⁻²¹ While TBI preceding PNES are mild (mTBI) in a majority of cases, moderate and severe TBI can also be associated with PNES.⁶ The fact that mTBI is much more common in PNES than moderate or severe TBI could lead to the perception that mTBI is somehow more pathogenic in relation to PNES compared to more severe trauma; however, this distribution more likely reflects the overall incidence of mild, moderate and severe TBI, with 80-90% of TBI in the general population categorized as mTBI.²² The latency between TBI and PNES onset varies greatly. In the two studies on adults which report this information (sample size 33 and 37) 81-89% patients developed PNES within the first year after TBI.^{3,4}

The epidemiology of TBI-PNES co-occurrence should be set against the general prevalence of both disorders. A meta-analytic study has calculated that 12% of the general population in developed countries reports a history of TBI (mostly mild), with men affected twice as often as women.²³ The incidence of hospital-treated cases of mTBI is about 100-300/100,000.²⁴ PNES is diagnosed at a rate of 1.4-4.9/100,000/year, has an estimated prevalence of 2-33/100,000, and shows a clear preponderance of women.²⁵ Clearly, when analysing the co-occurrence of TBI and PNES, these epidemiological data indicate that part of the correlation might be coincidental rather than causal, and that any causal relationship would be far from universal.

Table 1: History of TBI in patients with PNES

Year	PNES, n	TBI-Hx, n (%)	average age (range or \pm SD), gender of TBI-Hx subgroup	Latency	mild TBI	Ref.
1993	27	18 (67%)	-	-	-	15
1993	93	15 (16%)	-	-	-	16
1998	157	37 (24%)	34 y (15-56), 68% women	89% within 1 y	78%	3
1998	102	33 (32%)	34 y (17-57), 48% women	81% within 1 y	91%	4
1999	100	52 (52%)	-	-	-	14
2000	16 ^a	7 (44%)	10.5 y (5-18), 31% girls	71% within 7 mo	57%	13
2001	40	33 (82.5%)	-	-	-	17
2002	23	20 (83%)	-	-	100%	12
2004	-	34 ^b	36 y, 53% women	-	0% ^b	6
2005	63	16 (26%)	-	-	-	18
2006	21	10 (48)	-	-	-	19
2010	64	13 (20%)	-	-	"most"	8
2010	44	13 (44%)	-	-	-	20
2013	92	41 (45%)	38.9 y (\pm 12.6), 66% women	-	73%	11
2014	324	119 (37%)	37.3 y (\pm 12.2), 71% women	-	-	10
2015	77	38 (57%)	49 y, 19% women ^c	-	87%	5
2017	51	26 (51%)	-	-	-	7
2017	49	13 (27%)	-	-	-	9
2017	17	10 (59%)	-	-	-	21

^a Paediatric population^b Study recruited only patients with moderate or severe TBI preceding seizure disorder^c Study from veteran's hospital

PNES, psychogenic nonepileptic seizures; TBI, traumatic brain injury; TBI-Hx, positive history of head injury; mTBI, mild traumatic brain injury; y, years; mo, months; SD, standard deviation

Table 2: Potential aetiological factors underlying PNES and antecedent TBI

Non-lesional phenomena	Possible effect on mTBI-PNES correlation
Recall bias and misattribution	promotes illness belief that can perpetuate dysfunction
Emotional trauma	affective stressor can induce long-term changes in stress-responsivity and metacognitive function akin to the development of posttraumatic stress disorder (PTSD)
Maladaptive emotional learning	acute stress and concussive symptoms lead to aversively conditioned neuropsychiatric dysfunction ("seizure scaffold") that is adapted as a response to emotional, somatic or trauma-related cues
Lesional phenomena	
Axonal injury and alteration of global functional connectivity	disruption or destabilization of metacognitive faculties such as cognitive inhibition and self-awareness which predisposes to dissociation

PNES, psychogenic nonepileptic seizures; mTBI, mild traumatic brain injury

Non-lesional effects of head trauma

Most authors of correlational studies on TBI and PNES have offered only restrained speculations on causality. While a structural component in PNES development after TBI has been entertained,²⁶ psychological phenomena and confounders are much more commonly cited (Table 2 and Fig. 1).^{3,4} One possible factor is recall bias.²⁶ PNES patients are often given imprecise explanations of their attacks and can find themselves actively searching for conceivable causes of their symptoms. Inconsistent definitions of what constitutes TBI and concussion²⁷ can further add to this, with harmless “knocks to the head” being reinterpreted as disease triggers. In one study, 39% of patients ascribed their PNES to antecedent TBI independent of (and usually contrary to) their physician’s assessment.²⁶ This might especially be the case in the veteran community, where the causal association between TBI and (epileptic) seizures is common knowledge. In a disorder where symptom expectations play a key role (see below), etiological misattributions of this sort may gain secondary importance by promoting relevant illness beliefs.

TBI is a significant stressor that could precipitate PNES due to its emotional impact alone following established models of psychological trauma and dissociation.^{4,28} High rates of traumatic experiences in general and physical injury in particular have been associated with PNES.²⁹ Mild TBI can induce high levels of stress and anxiety,³⁰ so it might trigger neuropsychiatric dysfunction in predisposed individuals on an emotional level as would any other traumatic event. In line with this explanation, a study on veterans with seizures attributed to head trauma revealed that 57% of all patients with PNES after TBI had documented PTSD.⁵ Since PTSD has been identified as a strong independent predictor of PNES,²⁹ mTBI might be one step removed from PNES or entirely unrelated. Highlighting the role of emotional stress in presumed head injury sequelae, a prospective study on 175 trauma patients found that acute post-concussion syndrome is as likely to occur patients with mTBI as it is in non-head-injured trauma controls.³¹

Accidental head trauma combines a highly stressful situation (e.g. sports competition, vehicle accident, brawl, injury- and treatment-related pain and dysfunction) with acute neuropsychiatric impairments in motor control, sensory faculties, consciousness or memory. In this context of acute stress, maladaptive emotional learning (especially aversive conditioning) combined with affect-biased attentional prioritization³² can thus result in symptom modelling: a repertoire of dysfunction or “seizure scaffold” is coupled with an abnormal stress response to perceived threats.^{1,33} Put in computational terms, the expectation of seizure-like dysfunction, both consciously due to illness beliefs and unconsciously in response to conditioned emotional and somatic cues, is allotted such disproportionately high confidence (or “precision”), that it can override through active inference contradicting perceptual information and produce symptoms.³⁴ Such maladaptive behavioural cascades have been proposed to underlie other functional neurological disorders such as functional

dizziness³⁵ and sensory and motor conversion disorder.^{34,36} In a study of 50 patients with functional movement disorders, 80% reported a precipitating peripheral trauma and 39% of those retrospectively fulfilled the criteria for a panic attack in association with the trauma.³⁷ In a study of 869 cases of motor and sensory conversion disorder, 37% were preceded by some kind of physical injury.³⁸ Lastly, in a study of 100 consecutive patients with motor functional neurological disorders (PNES, functional weakness, functional movement disorders and mixed presentations), 42% of all patients reported antecedent head trauma, with a higher incidence in PNES and functional movement disorders patients than in functional weakness.⁷ These observations would support a cognitively and emotionally mediated connection between antecedent head trauma and PNES, independent of any structural neuronal dysfunction.

The pathology of mTBI

Mild TBI results from an external force to the head that generates intracranial pressure gradients through the inertia of the brain inside the skull.²² This can lead to shearing and strain forces which in turn cause microtubule disruption and primary and secondary damage to axons. Some neuropathological investigations have described multifocal axonal injury (typical for moderate and severe TBI) after mTBI, but confounders remain difficult to control for. Routine clinical neuroimaging is normal after mTBI, but axonal dysfunction and network connectivity alterations can be detected and quantified using diffusor tensor imaging (DTI).^{39,40} Even though mediators or confounders such as opiate use or anxiety disorders might be part of the causal chain linking TBI with brain connectivity alterations, the basic association itself has been well established. Furthermore, the degree of white matter abnormalities detected on DTI correlates well with cognitive performance following mTBI.³⁹ Several studies on patients with mTBI using tractography show that frontal association pathways are primarily (though not exclusively) affected. It is beyond the scope of this article to review the potential neuropsychiatric implications of each commonly affected tract, but the uncinate fasciculus, which is affected in 29% of patients after mTBI,⁴¹ will be discussed as an example. It connects anterior temporal structures such as the amygdala to prefrontal cortices and has been directly implicated in psychiatric disorders.⁴² Interestingly, in a study of moderate and severe TBI in children, DTI-measured injury to the uncinate fasciculus reliably predicted emotional and behavioural dysregulation.⁴³ Since the uncinate fasciculus undergoes developmental maturation well into the third decade of life⁴² post-concussive development of psychiatric disorders could in part be connected to its vulnerability and microstructural damage in both children and adults.

Another way to visualize the neurophysiological consequences of disruptions of long-range association fibres after mTBI is through functional magnetic resonance imaging (fMRI).⁴⁴ Studies using resting-state protocols have repeatedly demonstrated reduced functional connectivity of the

default mode network (DMN), which is closely associated with self-reflective mental activity, after mTBI.⁴⁴ Albeit mostly in subjects with moderate to severe TBI, Ham and colleagues have convincingly shown impairments of self-awareness to result from breakdown of functional interactions within the DMN, and not from focal posttraumatic lesions.⁴⁵ The same group has also shown DMN dysfunction after TBI to correlate with inefficient inhibitory cognitive control as assessed in a Stroop task.⁴⁶ Lastly, both electroencephalography (EEG) and magnetoencephalography (MEG) studies have shown disruptions of functional connectivity after mTBI.⁴⁷⁻⁴⁹ In conclusion, even though mTBI may often leave no traces on routine structural MRI, the disruption of functional network connectivity due to injury to long-range association fibres can have direct short- and medium-term effects on metacognitive functioning.

Dysconnectivity in PNES

To assess the possibility of a structural predisposition to PNES after mTBI, a pathophysiological commonality has to be demonstrated. While current conceptual frameworks for the development of PNES are supported by a wealth of clinical and psychometric data,^{1,28} empirical advances in search of the neurobiological underpinnings are still discordant.² Furthermore, the heterogeneity of PNES and their association with different neurological and psychiatric conditions makes a unique focal neural dysfunction unlikely. Still, PNES are dissociative events that invariably entail certain (meta-)cognitive deficits such as loss of inhibitory control, reduced sense of agency and emotion dysregulation,¹ which could be assumed to be related to distinct neurophysiological disruptions.

Several groups have investigated PNES pathology with structural and functional MRI. Using DTI, Hernando and colleagues showed a rightward asymmetry of uncinate fasciculus streamlines in PNES patients that was not seen in age- and sex-matched healthy controls.⁵⁰ Interestingly, age at PNES onset was correlated with asymmetry indices of fractional anisotropy. Since the average age at disease onset was 35 years, these correlations might be related to the prolonged maturation process of the uncinate fasciculus mentioned above. Another DTI study also found altered white matter structural connectivity in the left uncinate fasciculus, as well as the left corona radiata, left internal and external capsules, and left superior temporal gyrus.⁵¹ Ding and colleagues performed DTI and fMRI in PNES patients and found decreased network connectivity strength in both structural and functional connectivity networks compared to healthy controls.⁵² Using high-density EEG, Knyazeva and colleagues found that psychogenic seizure frequency was correlated with hypo-synchronization of prefrontal and parietal regions, which, as the authors note, are both integral parts of the DMN.⁵³ Several studies using various imaging and electrophysiological methods have assessed neural alterations in PNES populations, but their findings have yet to converge around a coherent model.⁵⁴⁻

⁶² While a unifying neural signature of PNES predisposition has yet to be defined, studies suggest that

network connectivity alterations could be a potentially defining feature. Metacognitive processes that rely on top-tier recursive large-scale networks might be most susceptible to such disruptions, facilitating dissociation, disinhibition, amnesia and loss of volition during moments of abnormal stress and neurohumoral activation.

Dissociogenic brain lesions: a hypothesis

With this in mind, a plausible pathophysiological link between mTBI-related disruptions of long-range connectivity and the development of PNES emerges which is additional rather than oppositional to the non-lesional mechanisms described earlier (Fig. 1). Conscious awareness of sensory input and behaviour requires global integration of brain signals, which is assumed to rely on finely tuned recursive long-range connectivity.⁶³ A disruption of global connectivity and metacognitive functioning following mTBI might predispose individuals to dissociative experiences in general and PNES in particular. Focusing on specific vulnerable pathways such as, for example, the uncinate fasciculus, further reveals possible biological links of this nature. Just as certain life events or psychiatric diseases can be seen as clear precursors or catalysts of PNES, certain brain insults could well be considered “dissociogenic” (in analogy to those considered “epileptogenic”). Multifocal injury and disruption of long-range axonal fibres after mTBI would be a good example for this.

Dissociative symptoms and disorders other than PNES can sometimes be observed following mTBI.⁶⁴ In a prospective study of 1116 acute trauma patients, the group that had suffered an mTBI ($n = 476$) were significantly more likely to report dissociative symptoms such as derealization, restricted affect, and reduced awareness.⁶⁵ In a similar study on 131 road traffic accident victims (aged 18-65 years, no known pre-existing psychological disorder) with ($n = 66$) and without ($n = 65$) mTBI, the mTBI-group reported significantly more dissociation since the trauma (excluding peri-traumatic dissociation) compared to non-mTBI participants.⁶⁶ Another study on 80 mTBI patients found exceptionally high scores on the Dissociative Experiences Scale a median of 24.5 weeks after injury, though potential biases in patient sampling limit the interpretability of this study.⁶⁷




Fig. 1. Potential pathways linking mTBI and PNES. Misattribution can foster illness beliefs that shape symptom expectations. Acute post-concussive symptoms can serve as a symptom model (“seizure scaffold”) that is aversively conditioned to emotional, somatic or trauma-related cues as part of a maladaptive stress response. The acute stress of mTBI, despite promoting peri-traumatic dissociation, could also lead to changes in metacognition through pathways described in PTSD. Lastly, mTBI-related axonal injury, which is thought to underlie some of the acute symptoms, can also alter global connectivity in ways that affect metacognition and promote dissociation. Abbreviations: PNES, psychogenic nonepileptic seizures; mTBI, mild traumatic brain injury; PTSD, posttraumatic stress disorder.

Testing and expanding the hypothesis

Previous attempts to localize specific brain lesions associated with dissociative phenomena besides PNES should be considered when formulating our hypothesis. Research into states and traits of dissociation in dissociative disorders (derealization and depersonalization disorder, dissociative identity disorder) and trauma-associated disorders (dissociative subtype of PTSD, borderline personality disorder) has relied mainly on functional task-related or resting-state imaging.⁶⁸ A long list of potentially relevant brain regions has thus been assorted, but questions of predisposition, causality, adaptation and compensation have yet to be elucidated. Since our hypothesis specifically regards brain injury, the “lesion method” of neuroanatomical localization seems more promising.⁶⁹ One mTBI study which comes close to addressing our hypothesis was performed by Niogi and colleagues and ingeniously applied a combination of trait-mapping and the lesion method using DTI.⁷⁰ In a first step, target cognitive domains (memory and attention) in healthy controls were found

to correlate significantly with specific white matter regions of interest (left hemisphere anterior corona radiata and uncinate fasciculus, respectively). In a second step, thus identified white matter tracts were studied in mTBI patients at least 1 month post-injury (range 1-53 months). Changes in microstructural white matter integrity (measured by fractional anisotropy) in the regions of interest indeed correlated with the respective cognitive faculties in the pathological range. There was no "cross-correlation" – uncinate fasciculus disruption did not correlate with attentional dysfunction and vice versa. This type of mechanistic studies using imaging and electrophysiology will be needed to test our hypothesis, in addition to large-scale epidemiological studies that more closely distinguish reported, suspected and confirmed TBI and register the presence and length of posttraumatic amnesia and loss of consciousness to allow for better classification.

What other types of dissociogenic brain insults could there be? New onset of PNES has been observed anecdotally after general neurosurgery⁷¹, as well as systematically after epilepsy surgery, affecting 2.4 – 8.8% of patients.⁷²⁻⁷⁴ Most of the preceding operations in epilepsy patients were complete or partial temporal lobe resections, in which the uncinate fasciculus and its limbic connections are routinely resected. In discussing the post-operative development of PNES, many reports cite the psychosocial stress of being seizure-free after surgery or the stress of surgical morbidity as potential psychological factors.^{72,73} However, the potentially deleterious effect of resective epilepsy surgery on global network connectivity has received little attention so far. Other examples of potentially dissociogenic pre-existing conditions in PNES patients include perinatal brain injury, anoxic brain damage and various epileptogenic lesions.

In conclusion, the idea of dissociogenic brain lesions predisposing to PNES does not negate or oppose established cognitive-behavioural mechanisms (Fig. 1), but rather emphasizes the inextricable concurrence of structural predispositions and functional disturbances in neuropsychiatric disorders. Recognizing and studying lesional effects on PNES pathology will enable a more comprehensive understanding of the disorder and help in devising treatment strategies. Furthermore, this train of thought can lead to a better conceptualization of other functional neurological disorders, as well as other lesional predispositions.

Declaration of interest statement:

Dr. Popkirov reports no disclosures relevant to this study. Dr. Carson gives independent testimony in court on a range of topics that include functional neurological symptoms. Dr. Stone runs a free non-profit self-help website www.neurosymptoms.org.

Acknowledgements

This work was supported in part by a scholarship for S. P. from the FoRUM Forschungsreferat (research office) of the Medical Faculty of the Ruhr-University Bochum, Germany.

References

1. Brown RJ, Reuber M. Towards an integrative theory of psychogenic non-epileptic seizures (PNES). *Clin Psychol Rev.* 2016b;47:55-70.
2. Mcsweeney M, Reuber M, Levita L. Neuroimaging studies in patients with psychogenic non-epileptic seizures: A systematic meta-review. *Neuroimage Clin* 2017;16:210-221.
3. Barry E, Krumholz A, Bergey GK, Chatha H, Alemayehu S, Grattan L. Nonepileptic posttraumatic seizures. *Epilepsia.* 1998 Apr;39(4):427-431.
4. Westbrook LE, Devinsky O, Geocadin R. Nonepileptic seizures after head injury. *Epilepsia.* 1998;39(9):978-982.
5. Salinsky M, Storzach D, Goy E, Evrard C. Traumatic brain injury and psychogenic seizures in veterans. *J Head Trauma Rehabil.* 2015;30(1):E65-70.
6. Hudak AM, Trivedi K, Harper CR, et al. Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury. *J Head Trauma Rehabil.* 2004;19(4):290-295.
7. Matin N, Young SS, Williams B, et al. Neuropsychiatric Associations With Gender, Illness Duration, Work Disability, and Motor Subtype in a U.S. Functional Neurological Disorders Clinic Population. *J Neuropsychiatry Clin Neurosci.* 2017 Fall;29(4):375-382.
8. An DM, Wu XT, Yan B, Mu J, Zhou D. Clinical features of psychogenic nonepileptic seizures: a study of 64 cases in southwest China. *Epilepsy Behav.* 2010;17(3):408-11.
9. Asadi-Pooya AA, Tinker J. Delay in diagnosis of psychogenic nonepileptic seizures in adults: A post hoc study. *Epilepsy Behav.* 2017;75:143-145.
10. Elliott JO, Charyton C. Biopsychosocial predictors of psychogenic non-epileptic seizures. *Epilepsy Res.* 2014;108(9):1543-1553.
11. LaFrance WC Jr, Deluca M, Machan JT, Fava JL. Traumatic brain injury and psychogenic nonepileptic seizures yield worse outcomes. *Epilepsia.* 2013;54(4):718-725.
12. Mogleby K, Blomhoff S, Malt UF, Dahlstrom A, Tauboll E, Gjerstad L. Psychiatric comorbidity and hostility in patients with psychogenic nonepileptic seizures compared with somatoform disorders and healthy controls. *Epilepsia.* 2002;43(2):193-198.
13. Pakalnis A, Paolicchi J. Psychogenic seizures after head injury in children. *J Child Neurol.* 2000;15(2):78-80.
14. Kalogjera-Sackellares D, Sackellares JC. Intellectual and neuropsychological features of patients with psychogenic pseudoseizures. *Psychiatry Res.* 1999;86(1):73-84.
15. Bowman ES. Etiology and clinical course of pseudoseizures. Relationship to trauma, depression, and dissociation. *Psychosomatics.* 1993;34(4):333-342.
16. Lancman ME, Brotherton TA, Asconapé JJ, Penry JK. Psychogenic seizures in adults: a longitudinal analysis. *Seizure.* 1993 Dec;2(4):281-6.

17. Sackellares DK, Sackellares JC. Impaired motor function in patients with psychogenic pseudoseizures. *Epilepsia*. 2001;42(12):1600-1606.
18. Kotsopoulos I, de Krom M, Kessels F, Lodder J, Troost J, Twellaar M, van Merode T, Knottnerus A. Incidence of epilepsy and predictive factors of epileptic and non-epileptic seizures. *Seizure*. 2005;14(3):175-182.
19. Cragar DE, Berry DT, Fakhoury TA, Cibula JE, Schmitt FA. Performance of patients with epilepsy or psychogenic non-epileptic seizures on four measures of effort. *Clin Neuropsychol*. 2006 Sep;20(3):552-66.
20. Baslet G, Roiko A, Prensky E. Heterogeneity in psychogenic nonepileptic seizures: understanding the role of psychiatric and neurological factors. *Epilepsy Behav*. 2010;17(2):236-241.
21. Erba G, Bianchi E, Giussani G, Langfitt J, Juersivich A, Beghi E. Patients' and caregivers' contributions for differentiating epileptic from psychogenic nonepileptic seizures. Value and limitations of self-reporting questionnaires: A pilot study. *Seizure*. 2017;53:66-71.
22. Blennow K, Brody DL, Kochanek PM, et al. Traumatic brain injuries. *Nat Rev Dis Primers*. 2016;2:16084.
23. Frost RB, Farrer TJ, Primosch M, Hedges DW. Prevalence of traumatic brain injury in the general adult population: a meta-analysis. *Neuroepidemiology*. 2013;40(3):154-159.
24. Cassidy JD, Carroll LJ, Peloso PM, et al. Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*. 2004;(43 Suppl):28-60.
25. Asadi-Pooya AA, Sperling MR. Epidemiology of psychogenic nonepileptic seizures. *Epilepsy Behav*. 2015;46:60-65.
26. Salinsky M, Parko K, Rutecki P, Boudreau E, Storzbach D. Attributing seizures to TBI: Validation of a brief patient questionnaire. *Epilepsy Behav*. 2016;57(Pt A):141-144.
27. Mayer AR, Quinn DK, Master CL. The spectrum of mild traumatic brain injury: A review. *Neurology*. 2017;89(6):623-632.
28. Brown RJ, Reuber M. Psychological and psychiatric aspects of psychogenic non-epileptic seizures (PNES): A systematic review. *Clin Psychol Rev*. 2016a;45:157-182.
29. Fisman A, Alves-Leon SV, Nunes RG, D'Andrea I, Figueira I. Traumatic events and posttraumatic stress disorder in patients with psychogenic nonepileptic seizures: a critical review. *Epilepsy Behav*. 2004;5(6):818-825.
30. Harvey AG, Bryant RA. Predictors of acute stress following mild traumatic brain injury. *Brain Inj*. 1998 Feb;12(2):147-154.
31. Meares S, Shores EA, Taylor AJ, et al. Mild traumatic brain injury does not predict acute postconcussion syndrome. *J Neurol Neurosurg Psychiatry*. 2008;79(3):300-306.
32. Ehlers MR, Todd RM. Genesis and Maintenance of Attentional Biases: The Role of the Locus Coeruleus-Noradrenaline System. *Neural Plast*. 2017;2017:6817349.
33. Bautista RE, Gonzales-Salazar W, Ochoa JG. Expanding the theory of symptom modeling in patents with psychogenic nonepileptic seizures. *Epilepsy Behav*. 2008;13(2):407-9.

34. Edwards MJ, Adams RA, Brown H, Pareés I, Friston KJ. A Bayesian account of “hysteria.” *Brain*. 2012;135(11):3495-3512.
35. Popkirov S, Staab JP, Stone J. Persistent postural-perceptual dizziness (PPPD): a common, characteristic and treatable cause of chronic dizziness. *Pract Neurol*. 2018;18(1):5-13.
36. Stone J, Carson A, Aditya H, et al. The role of physical injury in motor and sensory conversion symptoms: a systematic and narrative review. *J Psychosom Res*. 2009;66(5):383-390.
37. Pareés I, Kojovic M, Pires C, et al. Physical precipitating factors in functional movement disorders. *J Neurol Sci*. 2014;338(1-2):174-177.
38. Stone J, Warlow C, Sharpe M. Functional weakness: clues to mechanism from the nature of onset. *J Neurol Neurosurg Psychiatry*. 2012;83(1):67-69.
39. Niogi SN, Mukherjee P. Diffusion tensor imaging of mild traumatic brain injury. *J Head Trauma Rehabil*. 2010;25(4):241-255.
40. Mayer AR, Ling J, Mannell MV, et al. A prospective diffusion tensor imaging study in mild traumatic brain injury. 2010;74(8):643-650.
41. Niogi SN, Mukherjee P, Ghajar J, et al. Extent of microstructural white matter injury in postconcussive syndrome correlates with impaired cognitive reaction time: a 3T diffusion tensor imaging study of mild traumatic brain injury. *AJNR Am J Neuroradiol*. 2008;29(5):967-973.
42. Alnæs D, Kaufmann T, Doan NT, et al. Association of Heritable Cognitive Ability and Psychopathology With White Matter Properties in Children and Adolescents. *JAMA Psychiatry*. 2018 Jan 24. doi: 10.1001/jamapsychiatry.2017.4277. [Epub ahead of print]
43. Johnson CP, Juranek J, Kramer LA, Prasad MR, Swank PR, Ewing-Cobbs L. Predicting behavioral deficits in pediatric traumatic brain injury through uncinate fasciculus integrity. *J Int Neuropsychol Soc*. 2011;17(4):663-673.
44. Shin SS, Bales JW, Edward Dixon C, Hwang M. Structural imaging of mild traumatic brain injury may not be enough: overview of functional and metabolic imaging of mild traumatic brain injury. *Brain Imaging Behav*. 2017;11(2):591-610.
45. Ham TE, Bonnelle V, Hellyer P, et al. The neural basis of impaired self-awareness after traumatic brain injury. *Brain*. 2014;137(Pt 2):586-597.
46. Bonnelle V, Ham TE, Leech R, Kinnunen KM, Mehta MA, Greenwood RJ, Sharp DJ. Salience network integrity predicts default mode network function after traumatic brain injury. *Proc Natl Acad Sci U S A*. 2012;109(12):4690-4695.
47. Cao C, Slobounov S. Alteration of cortical functional connectivity as a result of traumatic brain injury revealed by graph theory, ICA, and sLORETA analyses of EEG signals. *IEEE Trans Neural Syst Rehabil Eng*. 2010;18(1):11-19.
48. Dimitriadis SI, Zouridakis G, Rezaie R, Babajani-Feremi A, Papanicolaou AC. Functional connectivity changes detected with magnetoencephalography after mild traumatic brain injury. *Neuroimage Clin*. 2015;9:519-531.
49. Kumar S, Rao SL, Chandramouli BA, Pillai SV. Reduction of functional brain connectivity in mild traumatic brain injury during working memory. *J Neurotrauma*. 2009;26(5):665-675.

50. Hernando KA, Szaflarski JP, Ver Hoef LW, Lee S, Allendorfer JB. Uncinate fasciculus connectivity in patients with psychogenic nonepileptic seizures: A preliminary diffusion tensor tractography study. *Epilepsy Behav.* 2015;45:68-73.
51. Lee S, Allendorfer JB, Gaston TE, et al. White matter diffusion abnormalities in patients with psychogenic non-epileptic seizures. *Brain Res.* 2015;1620:169-176.
52. Ding JR, An D, Liao W, et al. Altered functional and structural connectivity networks in psychogenic non-epileptic seizures. *PLoS One.* 2013;8(5):e63850.
53. Knyazeva MG, Jalili M, Frackowiak RS, Rossetti AO. Psychogenic seizures and frontal disconnection: EEG synchronisation study. *J Neurol Neurosurg Psychiatry.* 2011;82(5):505-511.
54. Barzegaran E, Carmeli C, Rossetti AO, Frackowiak RS, Knyazeva MG. Weakened functional connectivity in patients with psychogenic non-epileptic seizures (PNES) converges on basal ganglia. *J Neurol Neurosurg Psychiatry.* 2016;87(3):332-337.
55. Barzegaran E, Joudaki A, Jalili M, Rossetti AO, Frackowiak RS, Knyazeva MG. Properties of functional brain networks correlate with frequency of psychogenic non-epileptic seizures. *Front Hum Neurosci.* 2012;6:335.
56. Ding J, An D, Liao W, et al. Abnormal functional connectivity density in psychogenic non-epileptic seizures. *Epilepsy Res.* 2014;108(7):1184-1194.
57. Li R, Li Y, An D, Gong Q, Zhou D, Chen H. Altered regional activity and inter-regional functional connectivity in psychogenic non-epileptic seizures. *Sci Rep.* 2015;5:11635.
58. Li R, Liu K, Ma X, et al. Altered Functional Connectivity Patterns of the Insular Subregions in Psychogenic Nonepileptic Seizures. *Brain Topogr.* 2015;28(4):636-645.
59. Umesh S, Tikka SK, Goyal N, Sinha VK, Nizamie SH. Aberrant gamma band cortical sources and functional connectivity in adolescents with psychogenic non-epileptic seizures: A preliminary report. *Psychiatry Res.* 2017;247:51-54.
60. van der Kruis SJ, Bodde NM, Vaessen MJ, et al. Functional connectivity of dissociation in patients with psychogenic non-epileptic seizures. *J Neurol Neurosurg Psychiatry.* 2012;83(3):239-247.
61. van der Kruis SJ, Jagannathan SR, Bodde NM, et al. Resting-state networks and dissociation in psychogenic non-epileptic seizures. *J Psychiatr Res.* 2014;54:126-133.
62. Xue Q, Wang ZY, Xiong XC, Tian CY, Wang YP, Xu P. Altered brain connectivity in patients with psychogenic non-epileptic seizures: a scalp electroencephalography study. *J Int Med Res.* 2013;41(5):1682-1690.
63. Sergent C, Dehaene S. Neural processes underlying conscious perception: experimental findings and a global neuronal workspace framework. *J Physiol Paris.* 2004;98(4-6):374-384.
64. Cantagallo A, Grassi L, Della Sala S. Dissociative disorder after traumatic brain injury. *Brain Inj.* 1999;13(4):219-228.
65. Broomhall LG, Clark CR, McFarlane AC, et al. Early stage assessment and course of acute stress disorder after mild traumatic brain injury. *J Nerv Ment Dis.* 2009;197(3):178-181.
66. Jones C, Harvey AG, Brewin CR. Traumatic brain injury, dissociation, and posttraumatic stress disorder in road traffic accident survivors. *J Trauma Stress.* 2005;18(3):181-191.

67. Mooney G, Speed J. The association between mild traumatic brain injury and psychiatric conditions. *Brain Inj.* 2001;15(10):865-877.
68. Krause-Utz A, Frost R, Winter D, Elzinga BM. Dissociation and Alterations in Brain Function and Structure: Implications for Borderline Personality Disorder. *Curr Psychiatry Rep.* 2017;19(1):6.
69. Sierra M, Lopera F, Lambert MV, Phillips ML, David AS. Separating depersonalisation and derealisation: the relevance of the "lesion method". *J Neurol Neurosurg Psychiatry.* 2002;72(4):530-532.
70. Niogi SN, Mukherjee P, Ghajar J, et al. Structural dissociation of attentional control and memory in adults with and without mild traumatic brain injury. *Brain.* 2008;131(Pt 12):3209-3221.
71. Reuber M, Kral T, Kurthen M, Elger CE. New-onset psychogenic seizures after intracranial neurosurgery. *Acta Neurochir (Wien).* 2002;144(9):901-907; discussion 907.
72. Asadi-Pooya AA, Asadollahi M, Tinker J, Nei M, Sperling MR. Post-epilepsy surgery psychogenic nonepileptic seizures. *Epilepsia.* 2016;57(10):1691-1696.
73. Ney GC, Barr WB, Napolitano C, Decker R, Schaul N. New-onset psychogenic seizures after surgery for epilepsy. *Arch Neurol.* 1998;55(5):726-730.
74. Glosser G, Roberts D, Glosser DS. Nonepileptic seizures after resective epilepsy surgery. *Epilepsia.* 1999;40(12):1750-1754.